


DO-TH 09/06

FLAVOR VIOLATION IN SUSY

G. HILLER

Institut für Physik, Technische Universität Dortmund, D-44221 Dortmund, Germany

I review recent opportunities from flavor for b -physics and collider searches.

1 Introduction: Flavor physics

The starting point for flavor physics is that known matter comes in generations Ψ_i , $i = 1, 2, 3$, where each generation carries identical charges under the Standard Model (SM) gauge group. Within the SM, the generations are solely distinguished by the Yukawa interactions, which couple the fermions to the Higgs boson. The non-diagonal structure of the Yukawa matrices allows for quark mixing and flavor change under the weak interaction. The strength of the transition of (quark) flavor $i \rightarrow j$ is encoded in the CKM mixing matrix element V_{ji} . In matrix form:

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \sim \begin{pmatrix} 1 & \lambda & \lambda^3 \\ -\lambda & 1 & \lambda^2 \\ -\lambda^3 & -\lambda^2 & 1 \end{pmatrix}, \quad (1)$$

where the phenomenological parametrization in terms of the Wolfenstein parameter $\lambda \simeq 0.22$ reflects the strong hierarchy present in the quark mixing. The third generation (t, b) is decoupled from the first two at order λ^2 . Today, masses and mixings including CP-violation are known input, with still improving accuracy through experimental and theoretical efforts. The origin of the hierarchical quark mass pattern and mixing, however, remains unexplained within the SM.

Flavor changing neutral current (FCNC) processes are suppressed in the SM by several effects: they arise only at the loop level, they are suppressed by flavor mixing and, except for the top quark contribution, by small mass splittings "GIM-mechanism". Since existing FCNC data are consistent with the SM, a mechanism to control FCNCs is required for any SM extension.

There are several possibilities and combinations thereof: i There is no flavor suppression from contributions beyond the SM to FCNCs. They are small because the scale of New Physics




Figure 1: RG evolution of MFV coefficients within AMSB for $\tan\beta = 10$. Figure taken from Ref. 5.

(NP) is high and not connected to the scale of electroweak symmetry breaking, hence, not in reach of the Large Hadron Collider (LHC). Since much of our reasoning about extending the SM is driven by electroweak symmetry breaking, we will not consider this any further and assume a TeVish NP scale. *ii* There is no (significant) flavor suppression but we allow for contributions to cancel to reproduce the data. Since not every sensible observable is measured yet (with sufficient precision) there is the intriguing possibility that the SM breaks down in the next round of experiments^{1,2}. Especially promising here are rare decays and $B_s - \bar{B}_s$ -mixing. *iii* There is a flavor suppression in the NP contributions similar to the one of the SM, which is λ^n . Models with the same amount of flavor violation as the SM with the only source of flavor being the Yukawa matrices, are termed Minimal Flavor Violation (MFV)-type^{3,4}. We discuss solutions to the flavor problem and their signatures for the case of supersymmetry (SUSY) below.

2 Flavor in SUSY

The superpotential of the minimal supersymmetric standard model (MSSM) with unbroken R -parity, neglecting leptons, reads as $W_{MSSM} = Q_i(Y_U)_{ij}H_u\bar{U}_j + Q_i(Y_D)_{ij}H_d\bar{D}_j - \mu H_d H_u$. It contains no further sources of flavor violation than the SM, that is, the up and down quark Yukawas, Y_U and Y_D . Its supersymmetric contribution to the MSSM Lagrangian is therefore always MFV. Whether the full model is MFV, depends then on the breaking of supersymmetry.

Within MFV, the SUSY-breaking greatly simplifies⁴, here for $SU(2)$ -doublet masses, as⁵

$$(m_{\tilde{Q}}^2)^T = z_1^q \mathbf{1} + z_2^q Y_U Y_U^\dagger + z_3^q Y_D Y_D^\dagger + z_4^q (Y_U Y_U^\dagger)^2 + z_5^q (Y_D Y_D^\dagger)^2 + \dots, \quad (2)$$

up to terms involving higher powers of the Yukawas and analogous expressions for singlets and the trilinear terms. For $z_{i \neq 1} \equiv 0$ the soft terms are called flavor blind. Well-known MFV models are gauge and anomaly mediation, and by construction, mSUGRA and the CMSSM. The constrained flavor structure of the MFV-type MSSM yields generic predictions: Highly degenerate squarks of 1st and 2nd generation and a 3rd generation decoupled at order λ^2 .

Renormalization group (RG) running modifies all MFV coefficients z_i . In particular, a model flavor blind at the scale of SUSY breaking such as gauge mediation develops squark masses with non-trivial flavor structure at the weak scale⁶. While also being MFV, the behavior of anomaly mediated SUSY breaking (AMSB) is of marked difference: The amount of flavor violation decreases from the high to the weak scale, see Fig. 1, where we show ratios $|z_{i>1}/z_1|$ as a function of the scale. Moreover, for low $\tan\beta$, AMSB becomes exactly flavor blind in the quasi-infrared fixed point limit of the top Yukawa⁵.




Figure 2: Rare decays within mAMSB with or without flavor for either sign of μ . Left: $b \rightarrow s\gamma$, right: $B_s \rightarrow \mu\mu$.

3 Probing squark flavor

How to experimentally identify MFV and non-MFV variants ?

3.1 With b -physics

It is well-known that within MFV, $\mathcal{O}(1)$ effects are possible in rare processes if $\tan\beta$ is largish. With its UV insensitivity, AMSB is particularly predictive. Analytical expressions for the full flavor structure, that is, the z_i and the "mass insertions" $(\delta)_{ij}$ exist⁵. AMSB predictions for the branching ratios of $b \rightarrow s\gamma$ and $B_s \rightarrow \mu^+\mu^-$ decays are shown in Fig. 2. The leptonic mode will place additional constraints on $\tan\beta$ (depending on the gravitino mass $m_{3/2}$) for $\mu < 0$ once the upper bound on its branching ratio improves. Since in AMSB the squarks are generically heavy, the impact of a flavor universal shift to avoid tachyonic sleptons on squark flavor is small⁵.

Deviations from the SM can be more dramatic in non-MFV models. Key indicators are right-handed FCNCs, the presence of non-CKM CP-violation or the breakdown of relations between observables linked by CKM. Recently, a lot of theory interest has focussed on angular distributions in $B \rightarrow (K^* \rightarrow K\pi)\mu^+\mu^-$ ^{7,8}, also $B \rightarrow Kl^+l^-$ ⁹ decays. Opportunities arise from CP-asymmetries in semileptonic $B \rightarrow K^*$ and $B_s \rightarrow \Phi$ transitions, which are doubly-Cabibbo suppressed within MFV (and the SM) and sensitive to a multitude of NP couplings⁷: Four CP-asymmetries are CP-odd, hence, can be extracted without flavor tagging. This is advantageous for $\bar{B}_s, B_s \rightarrow (\Phi \rightarrow KK)\mu^+\mu^-$ decays. Three CP-asymmetries are odd under T_N , the naive T-transformation. They can be order one with NP even if the strong phases are small such as predicted in the framework of QCD factorization for low dilepton invariant mass.

3.2 At ATLAS/CMS: Long live the stop

It is an interesting question whether MFV and its CKM-like squark mixing can be measured at colliders. In MFV, mixing between the stop and first two generations is suppressed $(\tilde{m}_Q^2)_{23}/\tilde{m}^2 \sim y_b^2 V_{cb} V_{tb}^* \sim 10^{-5} \tan\beta^2$, see Eq. (2). Such a tiny coupling can nevertheless be probed if $\tilde{t} \rightarrow c\chi^0$ is the dominant decay of the lightest stop $\tilde{t}$ and its rate is sufficiently suppressed¹⁰. Then,

$$\tau_{\tilde{t}} \sim \text{ps} \left(\frac{m_{\tilde{t}}}{100 \text{ GeV}} \right) \left(\frac{0.03}{\Delta m/m_{\tilde{t}}} \right)^2 \left(\frac{10^{-5}}{Y} \right)^2, \quad \Delta m = m_{\tilde{t}} - m_{\chi^0}, \quad Y_{\text{MFV}} \sim y_b^2 V_{cb} V_{tb}^*, \quad (3)$$

the lifetime of the stop is long and yields a macroscopic decay length. The flavor diagonal decay of the stop into top can be forbidden kinematically by having the stop-neutralino splitting Δm below the top mass. For $\Delta m > m_b$, also the 4-body decays $\tilde{t} \rightarrow b\chi^0 l\nu$ open up. Approximately,

$$\frac{\Gamma(\tilde{t} \rightarrow b\chi^0 l\nu)}{\Gamma(\tilde{t} \rightarrow c\chi^0)} \approx \frac{g^6 |V_{tb}|^2}{2(4\pi)^4} \frac{(\Delta m - m_b)^8}{[Y(\Delta m)]^2 m_W^4 m_{\chi^+}^2}. \quad (4)$$




Figure 3: The $\tilde{t}c\tilde{\chi}^0$ coupling Y versus $\Delta m/m_{\tilde{t}}$. The stop has macroscopic decay length below the solid curve and decays dominantly through FCNCs to the left of the dark band. Figure adopted from Ref. 10.

Relevant regions of stop parameters are shown in Fig.3 for $m_{\tilde{t}} = 100$ GeV. The stop has decay length $\beta\gamma\tau_{\tilde{t}} > 0.1$ mm and dominant FCNC decay in the light shaded (blue) region. Experimental constraints exist for larger values of Δm ¹¹. Observation of a long-lived stop would strongly support MFV since other scenarios have generically much larger $\tilde{t}c\tilde{\chi}^0$ couplings Y ¹⁰. A light stop is ingredient of electroweak baryogenesis and supports coannihilation of the relic density¹²; it can be realized in hypercharged anomaly mediation¹³, or with large trilinear A -terms.

4 Summary

What are the flavor quantum numbers of the new particles and SM partners related to electroweak symmetry breaking ? Already strong constraints exist from K , D - and B -observables, supporting models with flavor suppression not too far from the one in the SM, perhaps even exactly as in the SM, called MFV. The LHCb experiment and possibly super flavor factories² will greatly contribute to clarifying this question, but also the ATLAS/CMS experiments, as the possibility to learn about squark flavor mixing from a stop decay length measurement shows.

Acknowledgments

G.H. is happy to thank her collaborators, and the organizers for the opportunity to come and speak at Moriond EW 09. The work reported here is supported in part by the Bundesministerium für Bildung und Forschung, Berlin-Bonn, and the German-Israeli-Foundation (G.I.F.).

References

1. M. Artuso *et al.*, Eur. Phys. J. C **57**, 309 (2008).
2. S. Hashimoto *et al.*, Letter of intent for KEK Super B Factory, KEK-REPORT-2004-4; J. L. Hewett *et al.*, arXiv:hep-ph/0503261; M. Bona *et al.*, arXiv:0709.0451 [hep-ex].
3. R. S. Chivukula and H. Georgi, Phys. Lett. B **188**, 99 (1987); A. Ali and D. London, Eur. Phys. J. C **9**, 687 (1999); A. J. Buras and R. Buras, Phys. Lett. B **501**, 223 (2001).
4. G. D'Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, Nucl. Phys. B **645**, 155 (2002).
5. B. C. Allanach, G. Hiller, D. R. T. Jones and P. Slavich, JHEP **04** (2009) 088.
6. P. Paradisi, M. Ratz, R. Schieren and C. Simonetto, Phys. Lett. B **668**, 202 (2008); G. Colangelo, E. Nikolidakis and C. Smith, Eur. Phys. J. C **59**, 75 (2009).
7. C. Bobeth, G. Hiller and G. Piranishvili, JHEP **0807** (2008) 106.
8. U. Egede, T. Hurth, J. Matias, M. Ramon and W. Reece, JHEP **0811** (2008) 032; W. Altmannshofer *et al.*, JHEP **0901** (2009) 019; A. Bharucha, *these proceedings*.
9. C. Bobeth, G. Hiller and G. Piranishvili, JHEP **0712** (2007) 040.
10. G. Hiller and Y. Nir, JHEP **0803** (2008) 046.
11. V. M. Abazov *et al.* [D0 Collaboration], Phys. Lett. B **665** (2008) 1.

12. C. Boehm, A. Djouadi and M. Drees, Phys. Rev. D **62**, 035012 (2000); C. Balazs, M. S. Carena and C. E. M. Wagner, Phys. Rev. D **70**, 015007 (2004).
13. R. Dermisek, H. Verlinde and L. T. Wang, Phys. Rev. Lett. **100**, 131804 (2008).